



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

325. Proposed by E. B. ESCOTT, University of Michigan, Ann Arbor, Michigan.

Integrate $\int \frac{4x^6 - a^3}{\sqrt{(x^6 - a^3)}} dx$.

326. Proposed by C. N. SCHMALL, New York City.

Prove $\int_0^\infty \frac{[(\tan^{-1}ax)^2 - (\tan^{-1}bx)^2]dx}{x} = \frac{1}{4}\pi^2 (\log a - \log b)$.

327. Proposed by RICHARD P. LOCHNER, 214 North 63rd Street, Philadelphia, Pennsylvania.

A hound is at the middle point of the side of a square field, and a fox is at an adjacent corner. How far will the hound run to catch the fox if the fox runs on the perimeter of the field and the hound runs directly towards the fox at all times, the hound running n times as fast as the fox? Where will the race end?

328. Proposed by M. E. GRABER, A. M., Heidelberg University, Tiffin, Ohio.

Prove that, $\frac{E_m}{\pi/2m} \int_0^{\pi/2m} [\sin a + \sin(\frac{\pi}{m} + a) + \dots \sin(\frac{m-1}{m}\pi + a)] da = \frac{2m E_m}{\pi}$.
[Sheldon and Hausmann: *Dynamo Electric Machinery*, Vol. I, p. 51.]

MECHANICS.

269. Proposed by F. H. SAFFORD, Ph. D., University of Pennsylvania.

Three jointed rods, lengths 5, 2, and 5 units, are suspended from two pins which are 6 units apart and on a horizontal line, forming a freely jointed quadrilateral with the shortest side at the bottom. At the two lower joints are weights of A and $2A$, respectively. Find the position of rest, the reactions along each bar, and the pressures on the pins.

270. Proposed by W. J. GREENSTREET, M. A., Editor, *The Mathematical Gazette*, Burghfield, England.

A cycloid has its base vertical. Find the line of quickest descent from the middle point of the base, and its approximate inclination to the horizon.

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

186. Proposed by H. PRIME, Boston, Massachusetts.

Show that $\frac{(n+1)(n+2)\dots(2n-2)}{(n-1)!}$ is an integer for all values of n .